Mini Nutritional Assessment
MNA®

Last name: First name:

Sex: Age: Weight, kg: Height, cm: Date:

Complete the screen by filling in the boxes with the appropriate numbers. Add the numbers for the screen. If score is 11 or less, continue with the assessment to gain a Malnutrition Indicator Score.

Screening

A Has food intake declined over the past 3 months due to loss of appetite, digestive problems, chewing or swallowing difficulties?
0 = severe decrease in food intake
1 = moderate decrease in food intake
2 = no decrease in food intake ☐

B Weight loss during the last 3 months
0 = weight loss greater than 3kg (6.6lbs)
1 = does not know
2 = weight loss between 1 and 3kg (2.2 and 6.6 lbs)
3 = no weight loss ☐

C Mobility
0 = bed or chair bound
1 = able to get out of bed / chair but does not go out
2 = goes out ☐

D Has suffered psychological stress or acute disease in the past 3 months?
0 = yes 2 = no ☐

E Neuropsychological problems
0 = severe dementia or depression
1 = mild dementia
2 = no psychological problems ☐

F Body Mass Index (BMI) (weight in kg) / (height in m^2)
0 = BMI less than 19
1 = BMI 19 to less than 21
2 = BMI 21 to less than 23
3 = BMI 23 or greater ☐

Screening score ☐☐
(subtotal max. 14 points)

12-14 points:	Normal nutritional status
8-11 points:	At risk of malnutrition
0-7 points:	Malnourished

For a more in-depth assessment, continue with questions G-R

Assessment

G Lives independently (not in nursing home or hospital)
1 = yes 0 = no ☐

H Takes more than 3 prescription drugs per day
0 = yes 1 = no ☐

I Pressure sores or skin ulcers
0 = yes 1 = no ☐

J How many full meals does the patient eat daily?
0 = 1 meal
1 = 2 meals
2 = 3 meals ☐

K Selected consumption markers for protein intake
- At least one serving of dairy products (milk, cheese, yoghurt) per day yes ☐ no ☐
- Two or more servings of legumes or eggs per week yes ☐ no ☐
- Meat, fish or poultry every day yes ☐ no ☐

0.0 = if 0 or 1 yes
0.5 = if 2 yes
1.0 = if 3 yes ☐.☐

L Consumes two or more servings of fruit or vegetables per day?
0 = no 1 = yes ☐

M How much fluid (water, juice, coffee, tea, milk...) is consumed per day?
0.0 = less than 3 cups
0.5 = 3 to 5 cups
1.0 = more than 5 cups ☐.☐

N Mode of feeding
0 = unable to eat without assistance
1 = self-fed with some difficulty
2 = self-fed without any problem ☐

O Self view of nutritional status
0 = views self as being malnourished
1 = is uncertain of nutritional state
2 = views self as having no nutritional problem ☐

P In comparison with other people of the same age, how does the patient consider his / her health status?
0.0 = not as good
0.5 = does not know
1.0 = as good
2.0 = better ☐.☐

Q Mid-arm circumference (MAC) in cm
0.0 = MAC less than 21
0.5 = MAC 21 to 22
1.0 = MAC 22 or greater ☐.☐

R Calf circumference (CC) in cm
0 = CC less than 31
1 = CC 31 or greater ☐

Assessment (max. 16 points) ☐☐.☐

Screening score ☐☐.☐

Total Assessment (max. 30 points) ☐☐.☐

Malnutrition Indicator Score

24 to 30 points	☐	normal nutritional status
17 to 23.5 points	☐	at risk of malnutrition
Less than 17 points	☐	malnourished

Ref. Vellas B, Villars H, Abellan G, et al. *Overview of MNA® - Its History and Challenges.* J Nut Health Aging 2006; 10: 456-465.
Rubenstein LZ, Harker JO, Salva A, Guigoz Y, Vellas B. Screening for Undernutrition in Geriatric Practice: *Developing the Short-Form Mini Nutritional Assessment (MNA-SF).* J. Geront 2001; 56A: M366-377.
Guigoz Y. The Mini-Nutritional Assessment (MNA®) *Review of the Literature – What does it tell us?* J Nutr Health Aging 2006; 10: 466-487.

® Société des Produits Nestlé, S.A., Vevey, Switzerland, Trademark Owners
© Nestlé, 1994, Revision 2006. N67200 12/99 10M

For more information: www.mna-elderly.com